

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

## Research Overview

## Recent Advances in the Generation of Chemical Diversity Libraries

Manoj C. Desai, Ronald N. Zuckermann, and Walter H. Moos  
Chiron Corporation, Emeryville, California

Strategy, Management and Health Policy				
Venture Capital Enabling Technology		Preclinical Development Toxicology, Formulation Drug Delivery, Pharmacokinetics	Clinical Development Phases I-III Regulatory, Quality, Manufacturing	Postmarketing Phase IV

**ABSTRACT** In recent years, screening in combination with a diverse compound collection has become a powerful method for discovering leads for the ever-increasing number of new biologically active peptides, proteins, receptors, and enzymes discovered continually. As a result, the rapid generation and screening of compound libraries (collections) have recently become important major tools in the search for novel lead structures. Diverse collections of compounds have been acquired by many strategies; these include (1) natural products from plants, fermentation, marine organisms, insect toxins, and ethnic pharmacotherapies; (2) recombinant randomized peptide libraries (often referred to as biological diversity); (3) multiple peptide synthesis; and (4) non-peptidic synthetic libraries. The present review provides an overview of the recent advances in the field of peptide and non-peptidic synthetic libraries. The progress made thus far is broadly divided into two categories: (1) Amide based libraries. These libraries share the concepts of the peptide library strategies; much of the referenced work thus refers to peptides, reflecting the bias of the literature to date. (2) Non-amide based libraries. This promising technology combines solid phase synthesis with classical organic synthesis to provide large numbers of compounds with desirable bioavailability and pharmacokinetics for screening. The basic premise behind the second approach is that the high affinity ligands, when identified, will be much more likely to become useful therapeutic agents than the compounds discovered from amide based libraries. Synthesizing small heterocyclic ring systems that use ligands of diverse biological activity via combinatorial strategies is a fast developing branch of medicinal chemistry. We are at an early state in the development of combinatorial chemistry. However, this dramatic convergence of technologies represents a fundamental advance in medicinal chemistry and promises to play a major role in future drug discovery efforts. © 1994 Wiley-Liss, Inc.

**Key Words:** molecular diversity, solid-phase synthesis, peptoid, combinatorial chemistry

## INTRODUCTION

Recent advances in molecular biology, pharmacology, and automation have allowed the testing of minute amounts of compound ( $\leq 1$  nanomoles) in a relatively short time in a receptor binding or an enzymatic assay of interest. To the major drug companies who have collected large numbers of compounds (usually  $\geq 100,000$ ) over the years via chemical laboratory synthesis, there has existed a competitive edge

in utilizing them for the discovery of new lead structures via screening. Many of the initial leads of recently reported bioactive compounds have been dis-

Received June 8, 1994; final version accepted June 19, 1994.

Address reprint requests to Manoj C. Desai, Chiron Corporation, 4560 Horton Street, Emeryville, CA 94608.

covered by random screening of these corporate compound inventories [Clozel et al., 1993; Hsu et al., 1991; Snider et al., 1991]. Thus, screening in combination with a diverse compound collection can be a powerful method for discovering leads for the ever-increasing number of new biologically active peptides, proteins, receptors, and enzymes discovered continually. In contrast, the biotech companies without any compound collections, find themselves at a competitive disadvantage in the discovery of leads via screening. Thus, it is no surprise that companies such as Affymax, Chiron, Houghten, and Selectide, to name a few, have taken the lead in the area of generating chemical diversity in the form of combinatorial libraries. As a result, the rapid generation and screening of compound libraries have recently become important major tools in the search for novel lead structures.

Diverse collections of compounds have been acquired by many strategies; these include (1) natural products from plants, fermentation, marine organisms, insect toxins, and ethnic pharmacotherapies [Hylands and Nisbet, 1991]; (2) recombinant randomized peptide libraries (often referred to as biological diversity) [Bull et al., 1992]; (3) multiple peptide synthesis [Jung and Beck-Sickinger, 1992] and (4) non-peptidic synthetic libraries [Pavia et al., 1993; Gallop et al., 1994; Gordon et al., 1994]. The range of diversity has extended in a major way beyond natural products only in the last decade. Progressive chemical and biological means for artificially generating diversity have now far outstripped the libraries of the past, with millions upon millions of compounds readily available through a variety of approaches. Recombinant systems such as bacteriophage excel in producing hundreds of millions of peptides from a block set of 20 standard amino acids (19 L-amino acids plus glycine); in contrast, the chemical synthesis strategies may be less persuasive in terms of numbers, but their diversity may be limitless because of the endless variety of building blocks possible in such an approach.

In compound libraries the compounds are synthesized and stored as mixtures; sometimes the compounds, however, are synthesized individually in pools. This feature permits the synthesis of large numbers of compounds in a relatively short time; also the screening time is considerably reduced. For these libraries to be useful for discovering lead structures, there are subtle issues that must be balanced against each other. The faster pace of synthesis is achieved by generating deliberate mixture of compounds per vial. Logically, the library can be synthesized at a faster pace by increasing the number of compounds per vial. However, this must be weighed against the time it

takes to deconvolute a mixture. Toward this objective ingenious detection protocols have been formulated [Chu et al., 1992; Kerr et al., 1993; Nikolaiev et al., 1993; Ohlmeyer et al., 1993]. The magnitude of mixing order also affects the individual concentration of components in a bio-assay; thus it has a direct effect on the activity of the lead structure that will be discovered. As one would expect, the higher the mixing order, the more potent will be the lead structure and statistically fewer will be discovered that bind to the target. Further complicating the matter is the identification of reliable chemistry that would allow the synthesis of medicinally interesting structures in reasonable amounts and rapid time from readily available starting materials. Other issues that have an impact on library strategy include automation of the process, the data handling capacity, the availability of key building blocks, the quantity of each compound synthesized, and the molecular weights of the oligomers, which may determine ultimate absorption, distribution, metabolism and excretion parameters. *Not surprisingly, a fruitful library strategy requires carefully balancing various opposing issues such that the probability of finding a lead structure for a given assay with desirable bioavailability and pharmacokinetics parameters is increased.*

Almost all library protocols described to date rely on solid-phase synthesis techniques. An advantage offered over solution based methods is that the reactions can be driven to completion by employing excess reagents; product isolation is accomplished simply by washing away excess reagents from support-bound material. In recent years, a variety of polymers have been described for synthesis; these include cellulose discs [Eichler et al., 1989a,b; Frank, 1992a,b, 1993; Frank and Doring, 1988], cellulose paper sheets [Gausepohl et al., 1992], cotton fabric [Eichler et al., 1991], Kel-F-gstyrene and polystyrene-1% divinylbenzene [Albericio et al., 1989], long-chain polystyrene-grafted polyethylene film [Berg et al., 1990a,b], and controlled-pore glass resin [Chong et al., 1992]. Nonetheless, the resin obtained from polymerization of polystyrene-1% divinylbenzene continues to be the support of choice.

In the last decade, Geysen et al. [1984], Houghten [1985], Lam et al. [1991], and others using variants of the solid phase peptide synthesis (SPPS) independently generated libraries having greater than tens of thousands of peptides. The combinatorial relationship between the length of an oligomer and the number of variants at each position made the synthesis of large libraries possible. An oligomer of length  $n$  with  $X$  variants at each position will theoretically give  $X^n$  compounds (Table 1).

TABLE 1. Relation Between X<sup>n</sup> and Number of Distinct Molecules

Number of amino acid residues (X <sup>n</sup> )	Number of peptides
2	400
3	8,000
4	160,000
5	3,200,000
6	64,000,000
7	1,280,000,000
8	25,600,000,000

With solid phase peptide synthesis, pioneered by Merrifield [1963] as the state of the art, libraries of peptides were synthesized and used in drug research. The drawback is that, unlike compound inventories at major drug companies, these libraries contain only biopolymers from natural amino acids, their enantiomers, or a small number of unnatural amino acids. The metabolic instability of these compounds, owing to proteolysis, and their poor absorption characteristics, often render them poor drug candidates.

The present review provides an overview of recent advances in the field leading to non-peptidic synthetic libraries. The progress made thus far is broadly divided into two categories: (1) Amide based libraries. These libraries share the concepts of peptide library strategies; much of the referenced work thus far refers to peptides, reflecting the bias of the literature to date. (2) Non-amide libraries. This promising technology combines solid phase synthesis with classical organic synthesis to provide large numbers of compounds with desirable bioavailability and pharmacokinetics for screening. The bias behind the second approach is that the high affinity ligands, when identified, may be more likely to become useful therapeutic agents than the compounds discovered from amide based libraries (whether this proves true or not remains to be seen). Table 2 lists the key contributors in the field of combinatorial chemistry.

#### AMIDE BASED LIBRARIES

##### Synthesis of Peptides: "Multi-Pin" Method

In the early 1980s, the field of chemically generated molecular diversity was born when Geysen demonstrated that peptides could be synthesized in a reusable form in numbers that are orders of magnitude greater than ever before [Geysen, 1987, 1991; Geysen et al., 1984].

In this method, peptides can be synthesized on polyacrylate-grafted polyethylene rods arranged in a standard 96-well microtiter plate format. The format

TABLE 2. Representative Companies Involved in Generating Combinatorial Libraries

Libraries	Representative companies
Carbamates	Affymax, Chiron
Carbohydrates	Glycomed, Ontogen
Natural products	Shaman, Xenova, MYCOsearch
Non-peptide amides (peptoids)	ArQule, Chiron, Houghten, PharmacoPeia, Selectide, Sphinx-Genesis
Oligonucleotides (biological)	Darwin, NeXagen, Evotech
Oligonucleotides (chemical)	Darwin, Gilead, ISIS
Peptides (biological)	Affymax, Chiron, Genentech, Ixsys, Optein, Protein Engineering
Peptides (chemical)	Affymax, Bristol-Myers Squibb, Chiron, Houghten, Lilly, PharmacoPeia, Selectide, Terrapin
Polyamines	Houghten
Scaffold-based	Chiron, Ontogen, Selective
Small molecules	ArQule, Chiron, Parke-Davis, PharmacoPeia
Spatially directed	Affymax, Chiron
Tagged libraries	Affymax, Chiron, PharmacoPeia

allows 96 individual peptides (or mixtures) to be synthesized simultaneously at the ends of the rods (Fig. 1). The technique utilizes conventional SPPS methods, readily available materials and instrumentation, and has been used, for example, to synthesize more than 3,000 peptides in a week. Moreover, peptides can be used in assays while remaining bound to the rods, allowing their re-use for 20 or more tests, or the peptides can be cleaved and then tested in solution-phase assays [Bray et al., 1990, 1991b; Geysen, 1990].

One cleavage method yields products that contain a diketopiperazine moiety at the C-terminus, thereby introducing novel structural diversity upon cleavage (Fig. 2) [Valerio et al., 1991]. Through the use of new linkers or handles for attachment to the solid support,  $\beta$ -alanine acid or  $\beta$ -alanine methyl amide C-termini can also be generated [Valerio et al., 1991]. Alternatively, support-bound peptides can be cleaved with ammonia vapor to afford the corresponding C-terminal primary amides [Bray et al., 1991a]. An amino acid indexer that facilitates multi-pin use has been described [Carter et al., 1992].

The multi-pin approach is ideally suited for preparing a complete set of all possible overlapping synthetic peptides of given length homologous to a peptide sequence of interest [Geysen et al., 1985]. The power of the multi-pin method in pharmaceutical research has been demonstrated recently in replacement net approaches to the determination of structure-activity relationships (SARs) in a series of potent hexapeptides that bind endothelin receptors [Spellmeyer et al., 1993a,b], and in studies of substance P




Figure 1. Schematic representation of a pin, derivatized for solid phase synthesis.

[Wang et al., 1993a,b]. A general discussion of systematic screening for bioactive peptides has been published [Maeji et al., 1991].

A recent discussion considers practical strategies for the synthesis and testing of large libraries, taking into account the number of monomers available, as well as various optimization methods [Geysen and Mason, 1993]. The combination of epitope mapping with other SAR methods, made possible by multi-pin or alternative multiple peptide synthesis methods, has been used to enhance tumor necrosis factor (TNF) activity [Rathjen and Aston, 1993]. It has also been suggested that peptide ligands, especially as selected from a vast library of random peptide sequences, may have potential in generating ligands for large-scale affinity chromatography [Baumbach and Hammond, 1992].

#### Synthesis of Peptides: "Tea-Bag Approach"

Among the first alternatives to multi-pin peptide synthesis was the tea-bag approach [Houghten, 1985], where small amounts of resins representing individual peptides are enclosed in porous polypropylene containers, which resemble tea bags (Fig. 3). Resin-containing tea bags are immersed in individual solutions of amino acids, and standard SPPS procedures are utilized. Deprotection and washing steps are accomplished by mixing the tea bags together in a reaction vessel, followed by their subsequent separation for the next coupling. Finally, peptides are cleaved in a multi-vessel HF apparatus [Houghten, 1986; Houghten et al., 1986]. This method affords significant amounts of fully characterizable, solution-phase peptides, which may yield more relevant data than solid-support bound peptides. Neither analysis nor separation is required for the identification of biologically active peptides, since the biological assay identifies the tea bags that contain active peptides.

A semi-automated process allows the simultaneous synthesis of up to 120 different peptides [Beck-Sickinger et al., 1991]. This method has been extended to a combinatorial approach [Houghten et al., 1991], which permits the rapid synthesis and evalua-

tion of millions of individual, unmodified, free peptides, in quantities useful for screening in solution. For example, from libraries theoretically containing more than one million oligopeptides, nanomolar potency opioid ligands were identified using radioreceptor assays, and the antigenic determinant of a peptide recognized by a monoclonal antibody (MAb 125-10F3) was determined using a competitive enzyme-linked immunosorbent assay (ELISA) [Houghten et al., 1992; Houghten and Dooley, 1993].

#### Synthesis of Peptoids: "Mixed Resin Method"

As a chemical alternative to natural biopolymers that would be pharmaceutically relevant, Simon et al. have developed "peptoids" [Simon et al., 1992, 1994]. Peptides can be thought of as polymers of glycine with side chains projecting from the  $\alpha$  carbon atoms; in N-substituted glycine (NSG) peptoids, the side chains are attached to the amide nitrogen. In these NSG peptoids the achiral, flexible backbone is equipped with functional groups spaced at intervals comparable to those in naturally occurring peptides (Fig. 4).

This provides a structurally flexible molecule with functionalities required for molecular recognition. NSG peptoids are synthesized via a "sub-monomer route" [Zuckermann and Banville, 1992; Zuckermann et al., 1992a,b,c,d]; the sub-monomer route avoids tedious synthesis and deprotection of N-terminal Fmoc-protected glycine building blocks. The construction of NSG peptoids proceeds from the C-terminal to the N-terminal end, with two repetitious steps; firstly the coupling of  $\alpha$ -haloacetic acid and secondly the displacement of halide with amine (Fig. 5).

The advantage of these new structures over peptides are many: (1) one would expect them to have improved metabolic stability to natural proteases because of N-substitution, which they do (unpublished data); (2) a greater diversity of side chain functionalities can be incorporated from the large number of commercially available and functionally diverse primary amines; (3) the achiral nature of the backbone simplifies synthetic problems, especially racemization; (4) a higher inherent flexibility provides the possibility of probing a larger conformational space; and (5) the synthetic protocol retains advantageous features of solid phase chemistry, namely, modular stepwise assembly from readily accessible monomers, as well as the capacity for automation [Zuckermann and Banville, 1992]. NSG peptoids are often synthesized via a mix and split method pioneered for peptide libraries, as a mixture of ~500 peptoids per pool. If biological activity is observed, a pool can be deconvoluted via a small number of synthetic cycles to individual active compounds [Simon et al., 1991].




Figure 2. Formation of diketopiperazine linked peptides, which allows screening in solution. Reagents: (a)  $\text{CF}_3\text{COOH}$ ; (b) base or buffer.




Figure 3. The "tea bag" method of multiple peptide synthesis.

Given that submonomer synthesis allows incorporation of any of over 1,000 readily available primary amines, in principle  $10^9$  trimers and  $10^{12}$  tetramers can be prepared. This vast combinatorial variety allows a paradigm to select a minimum number of amines to maximize diversity. Toward this objective, flower plots for thousands of amines have been developed; flower plots graphically represent sixteen properties (five finger print multidimensional scaling [MDS] parameters, five atom layer MDS dimensions, five topological index principle components and log P) for a given amine. These flower plots can aid in designing libraries, for example, selecting similar or dissimilar sets of amines for lead discovery and/or lead optimization [Simon et al., 1994]. By combining the sub-monomer synthesis of NSG peptoids using di-





Figure 4. Comparison of a peptide chain and an NSG peptoid chain.

verse amines, selected with the help of this type of statistical analysis, CHIR 2279 and CHIR 4531 were discovered to have high affinity for the  $\alpha$ -adrenergic



X=O or NH

Figure 5. Synthesis of NSC peptoids by the submonomer method. Reagents: (a) DIC, DMF; (b) R-NH<sub>2</sub>, DMSO, 25°C; (c) repeat steps a and b.

and  $\mu$ -opiate receptors, respectively (Fig. 6) [Zuckermann et al., 1994].

### "One Bead-One Compound"

Hruby, Lam, and co-workers at Selectide significantly extended SPPS by the application of selectively cleavable linkers [Lebl et al., 1994; Patek and Lebl, 1990, 1991a,b]. The Selectide process is based on: (1) synthesis of diverse compounds on polymeric beads, with a single structure on each particle, (2) screening of this library in either a binding assay based on an enzyme-linked or fluorescent tag or in standard solution-phase assays, and (3) determination of the structure on the "active" bead via sequencing peptides by Edman degradation or using mass spectroscopic techniques [Kocis et al., 1993; Lam et al., 1991, 1993; Lebl et al., 1993; Nikolaiev et al., 1993].

A key feature is that the structure of the peptide on a given bead is not tracked. Rather, it relies on determining the structure only if the synthesized peptide on the bead has biological activity of interest. The library is generated by mixed resin synthesis using 120  $\mu$ m diameter beads; a single bead with three linkers contains about 300 pmol of peptide. In this approach, a polymeric support is divided into equal portions for coupling to each individual amino acid, and later recombined to create mixtures. Uniform coupling is ensured, since competition between amino acids due to unequal coupling rates is eliminated. Individual oligomers (or polymers) are combined during washing and deprotection, and then separated into individual portions for the next coupling. Resulting peptides exhibit a purely statistical distribution of sequences. Through this combinatorial approach, a complete library of peptides can be prepared rapidly in approximately equimolar amounts.

The assay protocol takes advantage of the selective cleavage of branched linkers. Thus, on each bead three copies of a peptide are synthesized. The scheme (Fig. 7) for assaying peptide libraries involves: (1) dispensing the library at 500 beads/well, releasing  $\frac{1}{3}$  of each peptide and testing of 500 peptides per well in a biological assay; (2) on finding a particular well having

interesting activity, 500 beads are dispensed at one bead per well and activity is deconvoluted to individual peptides using another  $\frac{1}{3}$  of the peptide; and finally (3) the remaining  $\frac{1}{3}$  of the peptide on the "active" bead is released and its structure determined by analytical chemistry. Recently, Lebl and co-workers have extended this strategy for the generation of libraries containing amide bonds having non-natural building blocks [Lebl et al., 1994; Salmon et al., 1993].

### "Light Directed Chemical Synthesis"

The development of very large scale immobilized polymer synthesis (VLSIPS) resulted from advances in microelectronics, combining solid-phase synthesis with photolithography and miniaturization [Adams et al., 1994; Fodor et al., 1991; Gallop et al., 1994; Pirrung et al., 1992] (Fig. 8). In this technique, the use of photochemically labile protecting groups allows the light-directed, spatially addressable, parallel, chemical synthesis of thousands of oligomers at defined sites on a micromolar scale. Combinatorial masking strategies are used to generate large numbers of compounds in a small number of chemical steps. This technique allows the use of unnatural amino acids, nucleotides, and, in principle, any set of chemically compatible building blocks. The VLSIPS technology has been reviewed [Dower and Fodor, 1991; Gallop et al., 1994; Gordon et al., 1994].

### Additional Developments

In addition to these major contributors, Corvas has reported automated synthesis of peptide C-terminal aldehydes as an important class of transition-state analogues (Fig. 9) [Murphy et al., 1992]. This method relies on the protection of an aldehyde as a stereochemically stable semicarbazone. After the desired number of deprotection/coupling cycles are complete, the protected peptide semicarbazone is treated with aqueous acid/formaldehyde to regenerate the aldehyde, and then cleaved from the solid support.

A challenge in using combinatorial libraries is





Figure 6. Structures and affinities of NSC peptoids for  $\alpha 1$ -adrenergic (CHIR 2279) and  $\mu$ -opiate (CHIR 4531) receptors.




Figure 7. Scheme for discovery of active peptides from a "one peptide per bead" library.

the characterization of members of the library with desired properties. For this, some prefer to rely on iterative resynthesis, while others depend on analytical chemistry. Yet another solution, a "tagging" method, attempts to solve the structure elucidation problem by co-synthesizing a sequenceable tag that encodes a series of steps and reagents used in the synthesis of each library element. The tag and corresponding library element are associated by their attachment to a given resin bead. Once a library ele-

ment is selected, the procedure used to synthesize it can be read by sequencing the tag. Oligonucleotide [Brenner and Lerner, 1992; Needles et al., 1993] and oligopeptide [Kerr et al., 1993; Nikolaiev et al., 1993] tags have both been reported. Still and co-workers have devised an alternative method that uses a chemically inert, multiple tag labeling scheme, which attaches to the beads tagging molecules that encode both the step number and the chemical reagent used in that step. The array of tags used forms a binary record of the synthetic steps for each bead; tags can be read directly from a single bead by electron capture capillary gas chromatography [Borchardt and Still, 1994; Ohlmeyer et al., 1993].

The convergence of solid-phase synthesis methods with general building block strategies and pharmacophore-driven concepts such as NSC peptoids [Simon et al., 1992], a significant contribution of the last decade, is only now becoming more commonplace. In search of polypeptide-like substances with altered backbones, synthesis of vinyllogous polypeptides [Hagihara et al., 1992] and oligocarbamates [Cho et al., 1993; Simon et al., 1991] have been described (Fig. 10). Vinyllogous polypeptides consist of repeating units of extended amino acids that have an (*E*)-ethenyl unit inserted between the carbonyl carbon and  $C_{\alpha}$ . The key step in the synthesis of certain oligocarbamates involves the coupling of an amino group with *N*-protected *p*-nitrophenyl carbonate monomers with a variety of side chains (Fig. 11).

The less explored possibility of generating a library of compounds in solution is very appealing if compounds can be synthesized as mixtures in equal






Figure 8. Spatially addressable arrays of peptides can be synthesized on a glass surface using photolabel protecting groups (X).

amounts without side products and byproducts. A recent report of coupling an amine with a carboxylic acid with polymer bound 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide (P-EDC) holds considerable promise for this purpose [Desai and Stephens Stramiello, 1993]. The polymer tether enables the urea and N-acyl urea, if any, to remain on the polymer; the product therefore can be isolated simply by filtration and evaporation of the filtrate (Fig. 12).

### SYNTHESIS OF NON-AMIDE BASED LIBRARIES

The strategies discussed above produce structures that may have less than optimal bioavailability and pharmacokinetic parameters required for oral delivery of drugs. Many orally available drugs have heterocyclic structures, with MW < 500. The development of solid-phase synthesis of heterocyclic compounds is in a most critical stage for the drug discovery process: (1) the combinatorial synthesis of heterocyclic structures on solid phase has the ability to provide a large number of structures for in vitro screening in an assay of biological interest; and (2) the lead structure discovered from these libraries may require less extensive modification to produce suitable drug candidates.

The solid-phase synthesis of non-peptides is not unprecedented, although it is little known to organic chemists. Rapoport [Crowley and Rapoport, 1976] and Leznoff [Leznoff, 1978], are among the early pioneers to explore ambiguous limitations of non-peptide solid-phase chemistry. For example, Dieckmann and other condensation reactions were explored in the 1970s and before [Crowley and Rapoport, 1970, 1976]. The preparation of monoacetates of symmetrical diols [Fyles and Leznoff, 1976] and insect sex attractants [Fyles et al., 1978; Leznoff and Fyles, 1977] has also been accomplished using polymer-supported




Figure 9. Synthesis of peptide C-terminal aldehydes. Reagents: (a) Boc amino acid aldehyde/NaOAc reflux in ethanol; (b) MBHA resin/(benzotriazol-1-yloxy)tris(dimethylamino)-phosphonium hexafluorophosphate.




Figure 10. Comparison of portions of peptides, vinylogous polypeptides, and oligocarbamates.




Figure 11. Synthesis of oligocarbamates. Reagents: (a) 1, HOBT, diisopropylethylamine, NMP; (b) piperidine, NMP; (c) cleavage with TFA after capping with acetic anhydride.




Figure 12. Polymer bound 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (P-EDC).

organic synthesis [Leznoff, 1974, 1978]. The ability to produce compounds with drug characteristics should greatly enhance the drug discovery process. The focus has now been shifted to the synthesis of certain classes of compounds on solid phase and then utilizing the synthetic process for library production. Recent advances in this field are summarized below.

#### Syntheses of Benzodiazepines and Hydantoins

The ubiquitous 1,4-benzodiazepine nucleus is found in many drugs and ligands with diverse biolog-

ical activity. Most recently, it has been found in ligands for CCK-A and -B antagonists, tat antagonists, platelet-activating factor antagonists, GPIIb/IIIa inhibitors, reverse transcriptase inhibitors, and ras farnesyltransferase inhibitors [Bunin et al., 1994]. Thus, it is not surprising that 1,4-benzodiazepines were the first class of compounds for which combinatorial synthesis was described. Bunin and Ellman from Berkeley [Bunin and Ellman, 1992; Bunin et al., 1994] and DeWitt et al. from Parke-Davis [DeWitt et al., 1993] independently described the synthesis of benzodiazepine derivatives on polymer support. In Ellman's approach, 1,4-benzodiazepines are constructed on solid support from three components: aminobenzophenones, aminoacids, and alkylating agents (Fig. 13).

The 2-aminobenzophenone derivatives are first attached to the polystyrene solid support employing an acid cleavable linker 4-hydroxymethylphenoxyacetic acid through either an hydroxy or carboxylic acid functionality on one of the aromatic rings of the ami-




Figure 13. Synthesis of 1,4-benzodiazepine derivatives on solid support. Reagents: (a) 20% piperidine in DMF; (b) N-Fmoc amino acid fluoride, 4-methyl-2,6-di-*t*-butylpyridine; (c) 5% acetic acid in DMF; (d) lithiated 5-phenylmethyl-2-oxazolidone in THF,  $-78^{\circ}\text{C}$ , followed by alkylating agent and DMF; (e) TFA/ $\text{H}_2\text{O}/\text{Me}_2\text{S}$  (85:5:10).

nobenzophenone. Synthesis of the benzodiazepine derivatives on solid support proceeds by removal of the 9-fluorenylmethoxycarbonyl (Fmoc) protecting group from intermediate 2 by treatment with piperidine in DMF; this is followed by coupling the 2-aminobenzophenone to an  $\alpha$ -N-Fmoc amino acid fluoride. Deprotection of the Fmoc followed by exposure of the resulting free amine to 5% acetic acid in N-methylpyrrolidone provides the benzodiazepine ring system.

The Parke-Davis strategy relies (Fig. 14) on internal release for formation of the benzodiazepine ring system. The advantage of this strategy is that the compounds that do not "react" remain attached to the resin; this option provides final products in high purity. Through such a synthesis, 2–14 mg of 40 different benzodiazepines corresponding to 9–63% yield of each benzodiazepine were isolated with estimated purities typically >90%.

DeWitt et al. also described the synthesis of hydantoin, a primary nucleus in the anti-epileptic drug phenytoin, and the aldose reductase inhibitor sorbinil [Sarges et al., 1982]. The hydantoin synthesis begins by treating resin bound  $\alpha$ -amino acids with isocy-

anates to provide resin bound ureas. Subsequent cyclization and cleavage provides the hydantoin. This three-step protocol was utilized for the synthesis of 39 hydantoin in 4–81% yield [DeWitt et al., 1993] (Fig. 15).

#### Nitrile Oxide-Alkene/Alkyne [3 + 2] Cycloaddition Reactions

Kurth and co-workers reported a multi-step process for the synthesis of 2,5-disubstituted tetrahydrofurans via a tandem 1,3-dipolar cycloaddition/electrophilic cyclization sequence, thereby demonstrating the versatility of polymer supported methodology [Beebe et al., 1992]. The requisite nitro compound for the generation of nitrile oxide was prepared by the Henry reaction of nitromethane with polymer bound aldehyde. After protection of the hydroxyl moiety as a TMS ether, followed by phenylisocyanate mediated dehydration and subsequent 1,3-dipolar cycloaddition with 1,5-hexadiene (2–3-fold excess), this route provided polymer bound isoxazoline (Fig. 16). Finally, electrophilic cyclization of the isoxazoline with iodine monochloride at  $-78^{\circ}\text{C}$  gave the 2,5-disubstituted tetrahydrofurans.




Figure 14. General route for synthesis of benzodiazepines. Reagents: (a) 1,2-dichloroethane,  $\Delta$ ; (b) TFA/ $\Delta$ .




Figure 15. General route for the synthesis of hydantoins. Reagents: (a)  $R_3NCO$ , DMF, sonic bath, 6 h; (b) 6M aq. HCl, 85–100°C.




Figure 16. Polymer supported synthesis of 2,5-disubstituted tetrahydrofurans. Reagents: (a) 1. DMSO,  $NaHCO_3$ , 155°C, 6 h; 2.  $CH_3NO_2$ ,  $Et_3N$ , THF/EtOH, 14 h; 3. TMSCL,  $Et_3N$ , THF, 24 h; (b)  $PhNCO$ ,  $Et_3N$ , PhH, 80°C, sealed tube, 1,5-hexadiene, 4 days; (c) ICl,  $CH_2Cl_2$ , -78°C, 1.5 h.

Recently, Pei and Moos investigated the formation of isoxazoles and isoxazolines through [3+2] cycloaddition reaction of nitrile oxides with alkynes and alkenes [Pei and Moos, 1994]. In this strategy, NSG peptoids with allyl and alkyne side chains were synthesized; heating of such NSG peptoids in toluene with nitrile oxides, generated in situ by reacting nitro alkyl compounds with phenyl isocyanate and trimethylamine, afforded the isoxazoles and isoxazolines in >80% purity, respectively (Fig. 17). This "post-modification" of NSG peptoid chains has the potential to create greater diversity than might otherwise possible.

#### Synthesis of Oligosaccharides Using Polymer-Bound Glycals

Danishefsky and co-workers have re-examined the solid-phase synthesis of oligosaccharides by employing a newly developed stereospecific glycosylation strategy [Danishefsky et al., 1993]. This approach promises to ease some of the drudgery common to the synthesis and purification of oligosaccharides. The technique is applicable to synthesizing glycopeptides and glycosylated proteins, as well as oligosaccharides, and could possibly form the basis for an automated carbohydrate synthesizer. In this approach, glycals




Figure 17. [3+2] Cycloaddition reactions of nitrile oxides on solid phase. Reagents: (a)  $C_2H_5NO_2$ , PhNCO, TEA, toluene,  $100^\circ C$ ; (b) 20% TFA,  $CH_2Cl_2$ , room temperature.




Figure 18. Synthesis of oligosaccharides using polymer-bound glycals. Reagents: (a) Hunig's base and  $CH_2Cl_2$ ; (b) 3,3-dimethyldioxirane,  $CH_2Cl_2$ ; (c)  $I_2$ ,  $ZnCl_2$ , THF.

are attached to polystyrene co-polymer with a silyl ether bond, and are activated to function as glycosyl donors with 3,3-dimethyldioxirane. Glycosidation is performed by reaction with a solution-based acceptor. Excess acceptor and promoter are removed by rinsing after each coupling, and the desired oligosaccharides are cleaved from the polymer by addition of tetra-*n*-butylammonium fluoride. The average yield per coupling cycle consisting of epoxidation and glycosidation is ~70% (Fig. 18).

Synthesizing small heterocyclic ring systems that use ligands of diverse biological activity via combinatorial strategies is a fast developing branch of medicinal chemistry. We are at an early state in the development of combinatorial chemistry. However, this dramatic convergence of technologies represents a fundamental advance in medicinal chemistry and promises to play a major role in future drug discovery efforts.

## REFERENCES

- Adams CL, Kochersperger LM, Mortensen RB, Adwin LA, Holmes CP (1994): New methodologies in the preparation of peptide libraries. In Crabb, JW (ed): *Techniques in Protein Chemistry*, V. New York: Academic Press, Inc., pp 525-532.
- Albericio F, Ruiz-Gayo M, Pedrosa E, Giralt E (1989): Use of polystyrene-1% divinylbenzene and Kel-F-g-styrene for simultaneous synthesis of peptides. *React Polymer* 10:259-268.
- Baumbach GA, Hammond DJ (1992): Protein purification using affinity ligands deduced from peptide libraries. *BioPharm* 5:24-35.
- Beck-Sickinger AG, Duerr H, Jung G (1991): Semiautomated T-bag peptide synthesis using 9-fluorenylmethoxycarbonyl strategy and benzotriazol-1-yl-tetramethyluronium tetrafluoroborate activation. *Pep Res* 4:88-94.
- Beebe X, Schore NE, Kurth MJ (1992): Polymer supported synthesis of 2,5-disubstituted tetrahydrofurans. *J Am Chem Soc* 114:10061-10062.
- Berg RH, Almdal K, Pedersen WB, Holm A, Merrifield R (1990a): Peptide synthesis method and solid support for use in the method. World Patent 2,749.

- Berg RH, Almdal K, Perdersen WB, Holm A, Tam JP, Merrifield RB (1990b): A simple approach to rapid parallel synthesis of multiple peptide analogs. *Pept Chem Struct Biol Proc Am Pept Symp* 11:1036-1037.
- Borchardt A, Still WC (1994): Synthetic receptor binding elucidated with an encoded combinatorial library. *J Am Chem Soc* 116:373-374.
- Bray AM, Maeji NJ, Geysen HM (1990): The simultaneous multiple production of solution phase peptides: Assessment of the Geysen method of simultaneous peptide synthesis. *Tetrahedron Lett* 31:5811-5814.
- Bray AM, Maeji NJ, Jhingran AG, Valerio RM (1991a): Gas-phase cleavage of peptides from a solid support with ammonia vapour. Application in simultaneous multiple peptide synthesis. *Tetrahedron Lett* 32:6163-6166.
- Bray AM, Maeji NJ, Valerio RM, Campbell RA, Geysen HM (1991b): Direct cleavage of peptides from a solid support into aqueous buffer. Application in simultaneous multiple peptide synthesis. *J Org Chem* 56:6659-6666.
- Brenner S, Lerner RA (1992): Encoded combinatorial chemistry. *Proc Natl Acad Sci USA* 89:5381-5383.
- Bull AT, Goodfellow M, Slater JH (1992): Biodiversity as a source of innovation in biotechnology. *Annu Rev Microbiol* 46:219-252.
- Bunin BA, Ellman JA (1992): A general and expedient method for the solid-phase synthesis of 1,4-benzodiazepine derivatives. *J Am Chem Soc* 114:10997-10998.
- Bunin BA, Plunkett MJ, Ellman JA (1994): The combinatorial synthesis and chemical and biological evaluation of 1,4-benzodiazepine library. *Proc Natl Acad Sci USA* 91:4708-4712.
- Carter JM, VanAlburt S, Lee J, Lyon J, Deal C (1992): Shedding light on peptide synthesis. *Biotechnology* 10:509-513.
- Cho CY, Morna EJ, Cherry SR, Stephens JC, Fodor SPA, Adams CL, Sundaram A, Jacobs JW, Schultz PG (1993): An unnatural biopolymer. *Science* 261:1303-1305.
- Chong P, Sydor M, Wu E, Klein M (1992): Resin-bound synthetic peptides: Use as antigens and immunogens. *Mol Immunol* 29:443-446.
- Chu Y-H, Avila LZ, Biebuyck HA, Whitesides GM (1992): Use of affinity capillary electrophoresis to measure binding constants of ligands to proteins. *J Med Chem* 35:2915-2917.
- Clozel M, Breu V, Burri K, Cassal J-M, Fischli W, Gray GA, Hirth G, Loffler B-M, Muller M, Neidhart W, Ramuz H (1993): Pathophysiological role of endothelin receptor antagonist. *Nature* 365:759-761.
- Crowley JJ, Rapoport H (1970): Cyclization via solid phase synthesis. Unidirectional Dieckmann products from solid phase and benzyl triethylcarbinyl pinelates. *J Am Chem Soc* 92:6363-6365.
- Crowley JJ, Rapoport H (1976): Solid-phase organic synthesis: Novelty or fundamental concept? *Acc Chem Res* 9:135-144.
- Danishefsky SJ, McClure KF, Randolph JT, Ruggeri RB (1993): A strategy for the solid-phase synthesis of oligosaccharides. *Science* 260:1307-1309.
- Desai MC, Stephens Stramiello LM (1993): Polymer bound EDC (P-EDC): A convenient reagent for formation of an amide bond. *Tetrahedron Lett* 34:7685-7688.
- DeWitt SH, Kiely JS, Stankovic CJ, Schroeder MC, Reynolds Cody DM, Pavia MR (1993): "Diversomers": An approach to nonpeptide, nonoligomeric chemical diversity. *Proc Natl Acad Sci USA* 90:6909-6913.
- Dower WJ, Fodor SPA (1991): The search for molecular diversity (II): Recombinant and synthetic randomized peptide libraries. *Annu Rep Med Chem* 26:271-280.
- Eichler J, Beyermann M, Bienert M (1989a): Application of cellulose paper as support material in simultaneous solid-phase peptide synthesis. *Collect Czech Chem Commun* 54:1746-1752.
- Eichler J, Beyermann M, Bienert M, Lebl M (1989b): Simultaneous peptide synthesis using cellulose paper as support material. *Pept, Proc Eur Pept Symp*, 20th 205-207.
- Eichler J, Furkert J, Bienert M, Rohde W, Lebl M (1991): Multiple peptide synthesis on carriers: Elucidation and characterization of an antibody binding site of CRF. *Peptides 1990 Proc Eur Pept Symp* 21:156-157.
- Fodor SPA, Read JL, Pirrung MC, Stryer L, Lu AT, Solas D (1991): Light-directed, spatially addressable parallel chemical synthesis. *Science* 251:767-773.
- Frank R (1992a): Method and apparatus for the rapid synthesis of substrate-bound or free peptides or oligonucleotides, flat material synthesized in this way and the use of such material. *World Patent* 4,366.
- Frank R (1992b): Spot-synthesis: An easy technique for the positionally addressable, parallel chemical synthesis on a membrane support. *Tetrahedron Lett* 48:9217-9232.
- Frank R (1993): Strategies and techniques in simultaneous solid phase synthesis based on the segmentation of membrane type supports. *Bioorg Med Chem Lett* 3:425-430.
- Frank R, Doring R (1988): Simultaneous multiple peptide synthesis under continuous flow conditions on cellulose paper discs as segmental solid supports. *Tetrahedron Lett* 44:6031-6040.
- Fyles TM, Leznoff CC (1976): The use of polymer supports in organic synthesis. V. The preparation of monoacetates of symmetrical diols. *Can J Chem* 54:935-942.
- Fyles TM, Leznoff CC, Weatherston J (1978): Some solid-phase syntheses of the sex attractant of the spruce budworm,—trans-11-tetradecenol. *J Chem Ecol* 4:109-116.
- Gallop MBRW, Dower WJ, Fodor SPA, Gordon EM (1994): Applications of combinatorial technologies to drug discovery. 1. Background and peptide combinatorial libraries. *J Med Chem* 37:1233-1251.
- Gausepohl H, Boulin C, Kraft M, Frank RW (1992): Automated multiple peptide synthesis. *Pept Res* 5:315-320.
- Geysen HM (1987): Antigenically active amino acid sequences. *U.S. Patent* 4,708,871.
- Geysen HM (1990): Method for the use and synthesis of peptides. *World Patent* 9,395.
- Geysen HM (1991): Method of determining antigenically active amino acid sequences. *Eur Patent* 138,855.
- Geysen HM, Mason TJ (1993): Screening chemically synthesized peptide libraries for biologically relevant molecules. *Bioorg Med Chem Lett* 3:397-404.
- Geysen HM, Meloen R, Barteling S (1984): Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid. *Proc Natl Acad Sci USA* 81:3998-4002.
- Geysen HM, Barteling S, Meloen R (1985): Small peptides induce antibodies with a sequence and structural requirement for binding antigen comparable to antibodies raised against the native protein. *Proc Natl Acad Sci USA* 82:178-182.

- Gordon EM, Barrett RW, Dower WJ, Fodor SPA, Gallop MA (1994): Application of combinatorial technologies to drug discovery. 2. Combinatorial organic synthesis, library screening strategies, and future directions. *J Med Chem Soc* 37:1385-1401.
- Hagihara M, Anthony NJ, Stout TJ, Clardy J, Schreiber SL (1992): Vinylogous polypeptides: An alternative peptide backbone. *J Am Chem Soc* 114:6568-6570.
- Houghten RA (1985): General method for the rapid solid-phase synthesis of large numbers of peptides: Specificity of antigen-antibody interaction at the level of individual amino acids. *Proc Natl Acad Sci USA* 82:5131-5135.
- Houghten RA (1986): Means for sequential solid phase organic synthesis and methods using the same. U.S. Patent 4,631,211.
- Houghten RA, Dooley CT (1993): The use of synthetic peptide combinatorial libraries for the determination of peptide ligands in radio-receptor assays: Opioid peptides. *Bioorg Med Chem Lett* 3:405-412.
- Houghten RA, Bray MK, Degraw ST, Kirby CJ (1986): Simplified procedure for carrying out simultaneous hydrogen fluoride cleavage of protected peptide resins. *Int J Pept Pro Res* 27:673-678.
- Houghten RA, Pinilla C, Blondelle SE, Appel JR, Dooley CT, Cuervo JH (1991): Generation and use of synthetic peptide combinatorial libraries for basic research and drug discovery. *Nature* 354:84-86.
- Houghten RA, Appel JR, Blondelle SE, Cuervo JH, Dooley CT, Pinilla C (1992): The use of synthetic peptide combinatorial libraries for the identification of bioactive peptides. *Pept Res* 5:351-358.
- Hsu M-C, Schutt AD, Hooly M, Slice LW, Sherman MI, Richman DD, Potash MJ, Volsky DJ (1991): Inhibition of HIV replication in acute and chronic infections in vitro by a Tat antagonist. *Science* 254:1799-1802.
- Hylands PJ, Nisbet LJ (1991): The search for molecular diversity (I): Natural products. *Annu Rep Med Chem* 26:259-269.
- Jung G, Beck-Sickinger AG (1992): Multiple peptide synthesis methods and their applications. *Angew Chem Int Ed Engl* 31:367-383.
- Kerr JM, Banville SC, Zuckermann RN (1993): Encoded combinatorial peptide libraries containing non-natural amino acids. *J Am Chem Soc* 115:2529-2531.
- Kocis P, Krchnak V, Lebl M (1993): Symmetrical structure allowing the selective multiple release of a defined quantity of peptide from a single bead of polymeric support. *Tetrahedron Lett* 34:7251-7252.
- Lam KS, Salmon SE, Hersh EM, Hruby VJ, Kazmierski WM, Kanpp RJ (1991): A new type of synthetic peptide library for identifying ligand-binding activity. *Nature* 354:82-84.
- Lam KS, Hruby VJ, Lebl M, Knapp RJ, Kazmierski WM, Hersh EM, Salmon SE (1993): The chemical synthesis of large random peptide libraries and their use for the discovery of ligands for macromolecular acceptors. *Bioorg Med Chem Lett* 3:419-424.
- Lebl M, Patek M, Kocis P, Krchnak V, Hruby VJ, Salmon SE, Lam KS (1993): Multiple release of equimolar amounts of peptides from a polymeric carrier using orthogonal linkage-cleavage chemistry. *Int J Pept Pro Res* 41:201-203.
- Lebl M, Krchnak V, Safar P, Stierandova A, Spetov NF, Kocis P, Lam KS (1994): Construction and screening of libraries of peptide and non-peptide structures. In Crabb, JW (ed): *Techniques in Protein Chemistry*. V. New York: Academic Press, Inc., pp. 541-548.
- Leznoff CC (1974): The use of insoluble polymer supports in organic chemical synthesis. *Chem Soc Rev* 3:65-85.
- Leznoff CC (1978): The use of insoluble polymer supports in general organic synthesis. *Acc Chem Res* 11:327-33.
- Leznoff CC, Fyles TM (1977): The use of polymer supports in organic synthesis. VIII. Solid phase syntheses of insect sex attractants. *Can J Chem* 55:1143-1153.
- Maeji NJ, Bray AM, Valerio RM, Seldon MA, Wang J-X, Geysen HM (1991): Systematic screening for bioactive peptides. *Pept Res* 4:142-146.
- Merrifield RB (1963): Solid phase peptide synthesis I. The synthesis of a tetrapeptide. *J Am Chem Soc* 85:2149-2154.
- Murphy AM, Dagnino R Jr, Vallar PL, Trippe AJ, Sherman SL, Lumpkin RH, Tamura SY, Webb TR (1992): Automated synthesis of peptide C-terminal aldehydes. *J Am Chem Soc* 114:3156-3157.
- Needles MC, Jones DC, Tane EH, Heinkel GL, Kochersperger LM, Dower WJ, Barrett RW, Gallop MA (1993): Generation and screening of an oligonucleotide-encoded synthetic peptide library. *Proc Natl Acad Sci USA* 90:10700-10704.
- Nikolaiev V, Stierandova A, Krchnak V, Seligmann B, Lam KS, Salmon SE, Lebl M (1993): Peptide-encoding for structure determination of nonsequenceable polymers within libraries synthesized and tested on solid-phase supports. *Pept Res* 6:161-170.
- Ohlmeyer MHJ, Swanson RN, Dillard LW, Reader JC, Asouline C, Kobayashi R, Wigler M, Still WC (1993): Complex synthetic chemical libraries indexed with molecular tags. *Proc Natl Acad Sci USA* 90:10922-10926.
- Patek M, Lebl M (1990): A safety-catch type of amide protecting group. *Tetrahedron Lett* 31:5209-5212.
- Patek M, Lebl M (1991a): Safety-catch anchoring linkage for synthesis of peptide amides by Boc/Fmoc Strategy. *Tetrahedron Lett* 32:3891-3894.
- Patek M, Lebl M (1991b): "Safety-catch" protecting groups in peptide synthesis. *Collect Czech Chem Commun* 57:508-524.
- Pavia MR, Sawyer TK, Moos WH (1993): The generation of molecular diversity. *Bioorg Med Chem Lett* 3:387-396.
- Pei Y, Moos WH (1994): Post-modification of peptoid side chains: [3+2] cycloaddition of nitrile oxides with alkenes and alkynes on the solid-phase. *Tetrahedron Lett* (in press).
- Pirrung MC, Read JL, Fodor SPA, Stryer L (1992): Very large scale immobilized peptide synthesis. U.S. Patent 5,143,854.
- Rathjen DA, Aston R (1993): Selective enhancement of tumour necrosis factor activity: Mapping regions with monoclonal antibodies. *Bioorg Med Chem Lett* 3:457-462.
- Salmon SE, Lam KS, Lebl M, Kandola A, Khattri PS, Wade S, Patek M, Kocis P, Krchnak V, Thorpe D, Felder S (1993): Discovery of biologically active peptides in random libraries: Solution-phase testing after staged orthogonal release from resin beads. *Proc Natl Acad Sci USA* 90:11708-11712.
- Sarges R, Howard HR, Kelbaugh PR (1982): Synthesis of optically active spirohydantoins by asymmetric induction. Hydantoin formation from amino nitriles and chlorosulfonyl isocyanates. *J Org Chem* 47:4081.
- Simon R, Bartlett P, Santi D (1991): Modified peptide and peptide libraries with protease resistance, derivatives thereof and methods of producing and screening such. WO 91/19735.
- Simon RJ, Kania RS, Zuckermann RN, Huebner VD, Jewell DA,

- Banville S, Ng S, Wang L, Rosenberg S, Marlowe CK, Spellmeyer DC, Tan R, Frankel AD, Santi DV, Cohen FE, Bartlett PA (1992): Peptoids: A modular approach to drug discovery. *Proc Natl Acad Sci USA* 89:9367-9371.
- Simon RJ, Martin EJ, Miller SM, Zuckermann RN, Blaney JM, Moos WH (1994): Using peptoid libraries [oligo N-substituted glycines] for drug discovery. In Crabb, JW (ed): *Techniques in Protein Chemistry, V*. New York: Academic Press, Inc., pp. 533-539.
- Snider RM, Constantine JW, Lowe III JA, Longo KP, Lebel WS, Woody HA, Drozda SF, Desai MC, Vinick FJ, Spencer RW, Hess H (1991): A potent nonpeptide antagonist of the substance P (NK1) receptor. *Science* 251:435-437.
- Spellmeyer DC, Brown S, Stauber GB, Geysen HM, Valerio R (1993a): Endothelin receptor ligands. Multiple D-amino acid replacement net approach. *Bioorg Med Chem Lett* 3:1253-56.
- Spellmeyer DC, Brown S, Stauber GB, Geysen HM, Valerio R (1993b): Endothelin receptor ligands. Replacement net approach to SAR determination of potent hexapeptides. *Bioorg Med Chem Lett* 3:519-524.
- Valerio RM, Benstead M, Bray AM, Campbell RA, Maeji NJ (1991): Synthesis of solution phase peptide analogues using the multipin peptide synthesis method. *Anal Biochem* 197:168-177.
- Wang J, Bray AM, DiPasquale AJ, Maeji NJ, Geysen HM (1993a): Application of the multipin peptide synthesis technique for peptide receptor studies: Substance P as a model system. *Bioorg Med Chem Lett* 3:447-450.
- Wang J, DiPasquale AJ, Bray AM, Maeji NJ, Geysen HM (1993b): Study of conformation-dependent binding of substance P to central NK1 receptors by systematic D-amino acid replacements. *Bioorg Med Chem Lett* 3:451-456.
- Zuckermann RN, Banville SC (1992): Automated peptide-resin deprotection/cleavage by a robotic workstation. *Pept Res* 5:169-174.
- Zuckermann RN, Kerr JM, Kent SBH, Moos WH (1992a): Efficient method for the preparation of peptoids [oligo(N-substituted glycines)] by submonomer solid-phase synthesis. *J Am Chem Soc* 114:10646-10647.
- Zuckermann RN, Kerr JM, Siani MA, Banville SC (1992b): Design, construction and application of a fully automated equimolar peptide mixture synthesizer. *Int J Pept Prot Res* 40:497-506.
- Zuckermann RN, Kerr JM, Siani MA, Banville SC (1992c): Identification of highest-affinity ligands by selection from equimolar peptide mixtures generated by robotic synthesis. *Proc Natl Acad Sci USA* 89:4505-4509.
- Zuckermann RN, Siani MA, Banville SC (1992d): Control of the Zymate robot with an external computer: Construction of a multiple peptide synthesizer. *Lab Robotics Autom* 4:183-192.
- Zuckermann RN, Martin EJ, Spellmeyer DC, Stauber GB, Shoemaker KR, Kerr JM, Figliozzi CM, Goff DA, Siani MA, Simon RJ, Banville SC, Brown EC, Wang L, Richter LS, Moos WH (1994): Discovery of nanomolar ligands for 7-transmembrane G-protein coupled receptors from a diverse (N-substituted)glycine peptoid library. *J Med Chem* 37:2678-2685.